

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Frequency of the K303R Mutation Stratified by Nodal Status



	<u>WT ER</u>	<u>MUT ER</u>
Axillary N-	16	10 (38%)
Axillary N+	4	23 (85%)
	—	—

p=0.00062, Fisher's Exact Test


The Mutation is Associated with
Advanced Breast Cancer

Hypothesis: Proliferative Advantage is due to Hypersensitivity to Estrogen


Growth Assay



Transactivation MDA-231



Matrigel Invasion Assay - Estrogen Depleted




The K303R ERα Mutation is Frequently Present in Postmenopausal, ER-positive Patients

Neo-adjuvant study of 30 primary invasive breast cancers from UK^{***}:

Node-negative
Postmenopausal
Size > 2 cm
ERα-positive


10% WT

0 → 2w → 12w



90% Mut

0 → 2w → 12w




*** M. Dowsett

Why Find Only A to G Mutation in Tumors?

A to G = K303R
 A to T = K303M
 A to C = K303T

Transactivation Assay




E₂
 ERα


Only K303R is Hypersensitive

The ER α Mutation Exhibits Altered Binding to TIF-2 and AIB1 ER Co-activators


GST Pull-Down Assay



Altered Dynamics of K303R ER α on the pS2 Promoter



Mutant: increase in estrogen-induced occupancy on the promoter




Mutant: p300 appears to be retained in the complex


The K303R ER α Mutant is Resistant to MTA2 Co-Repressor Inhibition Of ER α Activity

Transactivation Assays

MCF-7



T47D





e K303R ER Mutant is Resistant to NCoR and BRCA1 Co-Repressor Inhibition of ER α Activity

HeLa

Transactivation Assays

T47D




The K303 ER α Site is Acetylated by p300

In Vitro Acetylation Assay

p300

autoacetylation



Control

GST-ER

GST-K303R/ER

GST

3, GATA-1 motif

R X K K

Acetylated by HATs
(p300 & P/CAF)

WT ER motif

R S K K

Acetylated by p300,
not P/CAF

Mutant 303 R

Acetylation by p300
is blocked

Wang et al., JBC. 2001

Are Receptor Acetylation and Phosphorylation Coupled Events?

Acetylation

PO₄

p53 or H3


Total *In Vitro* Kinase Assay:
GST-ERs Hinge #253-310

+


MCF-7 Lysate

+


γ P32



WT
K303R



↑ PO₄ of K303R ERα




ER Coomassie
blue staining

K303R ERα Mutation Generates an Efficient PKA & PAK1 Phosphorylation Site

Kinase	Substrate Specificity		
PKA	K/RXS (100%)	K/KXS (1%)	
PAK1	RXS (100%)	KXS (3%)	


Tuazon, et al. Biochem 1997

WT ER: KKNS₃₀₅
K303R: KRNS₃₀₅



ERα S305 Phosphorylation Reduces Receptor Acetylation


Transactivation Assay



Mutant mimicking phosphorylation at S305 increases estrogen sensitivity

Acetylation Assay: GST-ERs (Hinge #253-310)

↓
Acetylate with p300



* Acetylation
Coomassie blue stain

Other examples where phosphorylation blocks p300 acetylation:
Positive co-activator PC4, Forkhead TF FOXO 3a




FIG. 1



REPLACEMENT DRAWING

2/9




FIG. 2



3/9

N Dis.


G A T C



WT

TH

G A T C



VAR→

N Adj.

G A T C



VAR→

FIG. 3




FIG. 4





GST-wtER α		GST-K303RER α	
1/5 Input	GST	control	E ₂
			

FIG. 5



6/9




FIG. 6



7/9





FIG. 7




18 month old virgin mice-H&E

FIG. 8




9/9

NonTsg



K303R Tg



pH1b

18 month old virgin mice – pH1b IHC

FIG. 9